

CLDN8 Recombinant antibody

Cat:B35080D**Company:** HaoKebio**Uniprot ID:**P56748**Applications:** IHC:1:200-1:1000**Organism:**Rabbit

IHC-Polymer:1:800-1:4000

Species reactivity:Human Mouse Rat

IHC-TSA:1:1000-1:5000

Molecular Weight Calculation: 25 kDa

IF:1:50

Observed Molecular Weight: 25 kDa**Background:**

This gene encodes a member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This protein plays important roles in the paracellular cation barrier of the distal renal tubule, and in the paracellular barrier to prevent sodium back-leakage in distal colon. Differential expression of this gene has been observed in colorectal carcinoma and renal cell tumors, and along with claudin-7, is an immunohistochemical marker for the differential diagnosis of chromophobe renal cell carcinoma and renal oncocytoma.

Protein full name:

Claudin 8

Synonyms:

HEL-S-79; Claudin-8

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Membrane

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Source of Reagents:

发表[中文论文]请标注:CLDN8(B35080D)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:CLDN8(B35080D) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.